arXiv:hep-th/0505178v2 20 May 2005

The Large N Reduction in Matrix Quantum Mechanics — a Bridge between BFSS and IKKT —

Naoyuki Kawahara[1,2*] and Jun Nishimura[1,3†]
[1]High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801, Japan
[2]Department of Physics, Kyushu University, Fukuoka 812-8581, Japan
[3]Department of Particle and Nuclear Physics, School of High Energy Accelerator Science,
Graduate University for Advanced Studies (SOKENDAI), Tsukuba 305-0801, Japan

(Dated: May, 2005; preprint: KEK-TH-1018, KYUSHU-HET-84, hep-th/0505178)

The large N reduction is an equivalence between large N gauge theories and matrix models discovered by Eguchi and Kawai in the early 80s. In particular the continuum version of the quenched Eguchi-Kawai model may be useful in studying supersymmetric and/or chiral gauge theories nonperturbatively. We apply this idea to matrix quantum mechanics, which is relevant, for instance, to nonperturbative studies of the BFSS Matrix Theory, a conjectured nonperturbative definition of M-theory. In the bosonic case we present Monte Carlo results confirming the equivalence directly, and discuss a possible explanation based on the Schwinger-Dyson equations. In the supersymmetric case we argue that the equivalence holds as well although some care should be taken if the rotational symmetry is spontaneously broken. This equivalence provides an explicit relation between the BFSS model and the IKKT model, which may be used to translate results in one model to the other.

PACS numbers: 11.25.-w; 11.25.Sq

Introduction.— The large N reduction [1] states that large N gauge theories are equivalent to matrix models obtained by dimensionally reducing those theories to a point. Recent developments in the exact calculation of the low-energy superpotential in $\mathcal{N}=1$ supersymmetric gauge theories [2] may also be regarded as a particular application of the large N reduction [3]. We apply the original idea to matrix quantum mechanics, which is relevant, for instance, to nonperturbative studies of the BFSS Matrix Theory [4] proposed as a nonperturbative definition of M-theory. The same model may also be interpreted as a low-energy effective theory of D0-branes in type IIA superstring theory. (In the cases where fermions are absent, interesting results are already obtained by a direct lattice approach [5]. See Ref. [6] for a lattice formulation in the supersymmetric case.)

In Ref. [1] it was conjectured that the U(N) gauge theory on an infinite lattice is equivalent in the large N limit to a one-site model called the Eguchi-Kawai (EK) model [1]. This conjecture was based on the observation that the Schwinger-Dyson equations of the two theories coincide under the assumption that the U(1)^d symmetry $U_\mu \to e^{i\alpha_\mu}U_\mu$ of the EK model remains unbroken. The U(1)^d symmetry, however, turned out to be spontaneously broken in the weak coupling region [7, 8], and the quenched EK (QEK) model was proposed as a remedy [7]. The "quenched" eigenvalues of U_μ were interpreted later as lattice momenta [9], and this interpretation enabled an extension to non-gauge theories. The equivalence for the QEK model was confirmed both perturbatively [10] and nonperturbatively [7, 11].

The continuum version of the QEK (cQEK) model has been proposed in Ref. [12]. In spite of its potential usefulness in studying supersymmetric and/or chiral gauge theories, the EK equivalence for the cQEK model has not been checked directly so far.

The twisted EK model [13], which was proposed as an alternative way to cure the problem of the original EK model, has been interpreted recently as field theories on a noncommutative geometry [14]. However, this model is not useful for our purpose since the twisting procedure is applicable only to even space-time dimensions, and moreover the continuum version can be defined only in the strict $N=\infty$ limit [15]. As another recent development, it has been found that the breaking of the U(1)^d symmetry can be avoided by keeping the size of the lattice sufficiently large, and that the critical size in physical units becomes finite in the continuum limit [16].

In this letter we consider the cQEK model for matrix quantum mechanics, and discuss its EK equivalence. In fact the corresponding cQEK model is analogous to the IKKT model [17] proposed as a nonperturbative formulation of type IIB superstring theory. Based on this observation, we discuss an explicit relation between the BFSS model and the IKKT model.

The models.— We study a one-dimensional U(N) gauge theory, or matrix quantum mechanics (QM), with adjoint scalars $X_i(t)$ ($i=1,\cdots,d-1$), which are $N\times N$ Hermitian matrices. The action is given by

$$S_{\mathrm{mQM}} = \frac{1}{g^2}\int dt\, \mathrm{tr}\left\{\frac{1}{2}(DX_i(t))^2 - \frac{1}{4}[X_i(t),X_j(t)]^2\right\} , \tag{1}$$

where $DX_i = \frac{\partial}{\partial t}X_i - i\,[A, X_i]$ represents the covariant derivative with A being the 1d gauge field. The model can be obtained formally by the dimensional reduction of d-dimensional Yang-Mills theory to one dimension. In what follows we fix the gauge to $A(t)=0$. This model is UV finite and hence it does not require any UV regularization. As a result, the parameter g in (1) can always be scaled out by an appropriate rescaling of the matrices and the time coordinate t. We take $g = \frac{1}{\sqrt{N}}$, which turns out to be convenient when we discuss the large N limit.

Applying the quenching prescription [9, 12] to the matrix QM (1), we obtain the corresponding cQEK model

$$S_{\mathrm{cQEK},\Omega} = -N\,\epsilon\,\mathrm{tr}\left(\frac{1}{2}\,[\Omega, X_i]^2 + \frac{1}{4}\,[X_i, X_j]^2\right) \ , \qquad (2)$$

where X_i $(i = 1, \cdots, d-1)$ are $N \times N$ traceless Hermitian matrices and $\Omega \equiv \mathrm{diag}(\omega_1, \cdots, \omega_N)$ represents frequency variables to be integrated over the region $[-\frac{\Lambda}{2}, \frac{\Lambda}{2}]$, where $\Lambda \equiv \frac{2\pi}{\epsilon}$. (More precisely, the difference $\omega_i - \omega_j$ corresponds to the frequency.)

As U(N) invariant operators in the matrix QM (1), let us consider $\mathcal{O} = \frac{1}{N}\mathrm{tr}\, X_{i_1}(t_1) \cdots X_{i_n}(t_n)$ and the corresponding operator $\tilde{\mathcal{O}} = \frac{1}{N}\mathrm{tr}\, \tilde{X}_{i_1}(t_1) \cdots \tilde{X}_{i_n}(t_n)$ in the cQEK model (2), where $\tilde{X}_i(t) \equiv e^{i\Omega t}\, X_i\, e^{-i\Omega t}$. Then, according to the EK equivalence,

$$\lim_{N\to\infty} \langle \mathcal{O} \rangle_{\mathrm{mQM}} = \lim_{\Lambda\to\infty} \lim_{N\to\infty} \int d\Omega\, \langle \tilde{\mathcal{O}} \rangle_{\mathrm{cQEK},\Omega} \ , \qquad (3)$$

where $\langle \,\cdot\, \rangle_{\mathrm{mQM}}$ and $\langle \,\cdot\, \rangle_{\mathrm{cQEK},\Omega}$ represent the expectation values with respect to the actions (1) and (2), respectively, and $\int d\Omega \equiv \prod_{i=1}^{N} \int_{-\frac{\Lambda}{2}}^{\frac{\Lambda}{2}} \frac{d\omega_i}{\Lambda}$. The order of the two limits on the r.h.s. of (3) cannot be inverted.

Monte Carlo simulation.— We test the equivalence (3) directly by Monte Carlo simulation. In order to simulate the 1d model (1), we discretize the time "t", and obtain the action

$$\begin{aligned} S_{\mathrm{lat}} \ &= \ N a \left\{ \sum_{n=1}^{T-1} \frac{1}{2}\,\mathrm{tr}\left(\frac{X_i(n+1) - X_i(n)}{a} \right)^2 \right. \\ & \left. - \sum_{n=1}^{T} \frac{1}{4}\,\mathrm{tr}\,[X_i(n), X_j(n)]^2 \right\} \ , \qquad (4) \end{aligned}$$

where a represents the lattice spacing. The $N \times N$ Hermitian matrices $X_i(n)$ $(n = 1, \cdots, T)$ sit on each site, and we do *not* impose periodic boundary conditions. In order to retrieve the original model (1), we have to take the $a \to 0$ and $\tau \equiv a\,T \to \infty$ limits, where the order of the limits should not matter. The system is invariant under $X_i(n) \mapsto X_i(n) + \beta_i\,\mathbf{1}$, and we fix the corresponding zero mode by imposing the condition $\sum_{n=1}^{T} \mathrm{tr}\, X_i(n) = 0$.

Monte Carlo simulation of the models (4) and (2) can be performed by the heat bath algorithm [18]. For the cQEK model, we first take an average over the ensemble $\{X_i\}$ with the action (2) for each Ω, and then perform the averaging over Ω by using random numbers distributed uniformly within the region $[-\frac{\Lambda}{2}, \frac{\Lambda}{2}]$. Correspondingly there are two types of statistical errors. We estimate the total error by taking an average over the error of the first type and adding the one of the second type.

We focus on the $d = 4$ case in what follows. Let us first consider the observable $\mathcal{O}_1 = \frac{1}{NT} \sum_{n=1}^{T} \mathrm{tr}\,(X_i(n)^2)$ for the lattice model (4). In Fig. 1 we plot the results for $\langle \mathcal{O}_1 \rangle_{\mathrm{lat}}$ against a for $\tau = 5.0, 7.0, 10.0, 15.0$ at $N = 16$. We fit the results to $\langle \mathcal{O}_1 \rangle_{\mathrm{lat}} = C_1 + C_2\, a + C_3\, a^2$, from which we obtain the continuum limit.



FIG. 1: The observable $\langle \mathcal{O}_1 \rangle_{\mathrm{lat}}$ for the lattice model (4) with $N = 16$ is plotted against a for $\tau = 5.0, 7.0, 10.0, 15.0$. The lines represent fits to $\langle \mathcal{O}_1 \rangle_{\mathrm{lat}} = C_1 + C_2\, a + C_3\, a^2$.



FIG. 2: The observable $\langle \mathcal{O}_1 \rangle_{\mathrm{lat}}$ for the lattice model (4) after taking the $a \to 0$ and $\tau \to \infty$ limits are plotted against $\frac{1}{N^2}$. The straight line represents a fit to $C_6 + C_7\, N^{-2}$.

The results obtained in the continuum limit for various τ can be fitted to $\lim_{a\to 0} \langle \mathcal{O}_1 \rangle_{\mathrm{lat}} = C_4 + C_5\, \tau^{-1}$, from which we obtain the $\tau \to \infty$ limit. We redo this analysis for $N = 20, 32$. The results can be fitted to $C_6 + C_7\, N^{-2}$ as one can see from Fig. 2. This large N behavior is analogous to the one observed in the bosonic IKKT model [18]. Thus we can obtain the final result for the matrix QM (1).

Let us move on to the cQEK model (2). The corresponding observable is $\tilde{\mathcal{O}}_1 = \frac{1}{N}\mathrm{tr}\,(X_i^2)$. In Fig. 3 we plot the results $\langle \tilde{\mathcal{O}}_1 \rangle_{\mathrm{cQEK}}$ obtained for $N = 32, 48, 64$ with $\epsilon = 0.2, 0.25, 0.3, 0.5$. For fixed ϵ the large N behavior is given by $C_8 + C_9\, N^{-2}$. The coefficient C_9, however, diverges as $\frac{1}{\epsilon^2}$. This can be understood if we recall that the density of ω_i in the region $[-\frac{\Lambda}{2}, \frac{\Lambda}{2}]$ is $\rho = \frac{N}{\Lambda} = \frac{N\epsilon}{2\pi}$. Since finite N effects at small ϵ come mainly from the finiteness of ρ, any power of N should be associated with the same power of ϵ. This motivated us to plot the results against $\frac{1}{(N\epsilon)^2}$ in Fig. 3, where the results indeed lie on a single straight line for $\epsilon \leq 0.3$. Fitting all the data for $\epsilon \leq 0.3$, we obtain the final result for the cQEK model (2).



FIG. 3: The observable $\langle \tilde{\mathcal{O}}_1 \rangle_{\rm cQEK}$ for the cQEK model (2) is plotted against $\frac{1}{(N\epsilon)^2}$ for $\epsilon = 0.2$, 0.25, 0.3, 0.5. The straight line represents a fit to $\langle \tilde{\mathcal{O}}_1 \rangle_{\rm cQEK} = C_8 + C_{10}\,(N\epsilon)^{-2}$. The cross on the vertical axis (the error bar is almost invisible) represents the final result for the matrix QM (1).

We repeat this analysis for other observables and summarize the results in Table I, where we find good agreement within error bars. We consider this as a compelling evidence for the EK equivalence (3).

Schwinger-Dyson equations.— Let us discuss a possible explanation for the equivalence (3) based on the Schwinger-Dyson (SD) equations [1, 12, 19].

In the matrix QM (1) we consider

$$\langle {\rm tr}\,(\lambda^a X_{i_1}(t_1) \cdots X_{i_n}(t_n)) \rangle_{\rm mQM} \ ,$$

where λ^a represents a generator of the U(N) group. Changing the integration variables $X_i(t) \mapsto X_i(t) + \lambda^a \delta_{i,i_0} \delta(t - t_0)$, summing over a, using the identity $\sum_a (\lambda^a)_{ij} (\lambda^a)_{kl} = \delta_{il}\delta_{jk}$, and assuming the large N factorization property $\langle \mathcal{O}\mathcal{O}' \rangle_{\rm mQM} \simeq \langle \mathcal{O} \rangle_{\rm mQM} \langle \mathcal{O}' \rangle_{\rm mQM}$, we obtain a closed set of equations for the observables $\langle \mathcal{O} \rangle_{\rm mQM}$, which is the SD equations.

In the cQEK model we consider

$$\int d\Omega \, \langle {\rm tr}\,(e^{i\Omega t_0} \lambda^a e^{-i\Omega t_0} \tilde{X}_{i_1}(t_1) \cdots \tilde{X}_{i_n}(t_n)) \rangle_{{\rm cQEK},\Omega} \ .$$

Changing the integration variables $X_i \mapsto X_i + \frac{1}{\epsilon} \lambda^a \delta_{i,i_0}$, we proceed similarly to the matrix QM case assuming in addition that the Ω-integration factorizes as well. In this way we obtain a set of equations, which are the same as the SD equations for the matrix QM with the identification (3) except for the terms $(s = 1, \cdots, n)$

$$\frac{1}{\epsilon} \delta_{i_0,i_s} \int d\Omega \langle {\rm tr}\,(e^{i\Omega(t_s - t_0)} \tilde{X}_{i_1}(t_1) \cdots \tilde{X}_{i_{s-1}}(t_{s-1})) \rangle_{{\rm cQEK},\Omega}$$
$$\times \int d\Omega \langle {\rm tr}\,(\tilde{X}_{i_{s+1}}(t_{s+1}) \cdots \tilde{X}_{i_n}(t_n) e^{-i\Omega(t_s - t_0)}) \rangle_{{\rm cQEK},\Omega} \ ,$$

whose counterpart exists in the matrix QM only when $t_s = t_0$. If we neglect the fact that the integration region of Ω is bounded by Λ, the system is invariant under the shift $\Omega \mapsto \Omega + \alpha\,\mathbf{1}$ except for the factors $e^{\pm i\Omega(t_s - t_0)}$, which make the two Ω-integrations vanish separately for $t_s \neq t_0$.

Since the model we are studying is UV finite, we may expect that the integrand for the Ω-integration is determined by the ω_i's which are clustered in a finite region. Then the integration over the position of the cluster gives a factor of $f(t_s - t_0) = \int_{-\Lambda/2}^{\Lambda/2} \frac{d\omega}{\Lambda} e^{i\omega(t_s - t_0)}$, and we obtain the coefficient $\frac{1}{\epsilon} f(t_s - t_0)^2 \to \delta(t_s - t_0)$, which agrees with the corresponding one in the matrix QM (1) in the $\epsilon \to 0$ limit. Assuming that the SD equations have a unique solution, we obtain the equality (3).

Relation to the IKKT model.— The cQEK model (2) for the matrix QM has the action of the same form as the bosonic IKKT model [20] $S_{\rm bIKKT} = -\frac{N}{4} {\rm tr}\,[A_\mu, A_\nu]^2$ if we identify $A_i \equiv \epsilon^{1/4} X_i$ ($i \leq d-1$) and $A_d \equiv \epsilon^{1/4} \Omega$. The only difference is that A_d has to be quenched. In the bosonic IKKT model it is known that $\langle \frac{1}{N} {\rm tr}\,(A_\mu)^2 \rangle_{\rm bIKKT}$ is of O(1) [18], and the extent of the eigenvalue distribution is of O(1) in all directions. The quenching stretches the eigenvalue distribution of A_d to have an extent of O($\epsilon^{-3/4}$), and as a result the eigenvalue distribution of A_i ($i \leq d-1$) shrinks to O($\epsilon^{1/4}$) according to our results. This implies that, in spite of their formal similarity, the bosonic BFSS model and the bosonic IKKT model probe totally different regions of the configuration space.

How about the supersymmetric case? In $d = 4$, since the situation is qualitatively the same (no spontaneous breaking of the rotational symmetry and $\langle \frac{1}{N} {\rm tr}\,(A_\mu)^2 \rangle \sim$ O(1) [21]), all the statements for the bosonic case should equally apply.

In $d = 10$ (i.e., the IKKT model), there are certain evidences that the eigenvalue distribution of A_μ collapses dynamically to a four-dimensional hypersurface [22]. Let us assume further that the eigenvalue distribution extends to infinity in four directions at large N, and that the distribution is uniform in these directions, as expected if the IKKT model really describes our 4d space-

observable	matrix QM	cQEK
$\langle \frac{1}{N} {\rm tr}\, X_i^2 \rangle$	1.279(1)	1.28(1)
$\langle \frac{1}{N} {\rm tr}\, X_i^2 X_j^2 \rangle$	2.15(1)	2.11(5)
$\langle \frac{1}{N} {\rm tr}\, (X_i X_j)^2 \rangle$	1.23(2)	1.27(4)

TABLE I: Comparison of the Monte Carlo results obtained for the matrix QM (1) and the cQEK model (2).

time. When we quench the eigenvalues of A_{10}, the 4d hypersurface will arrange itself to have the required extent in the 10th direction. In this situation a vector in the 10th direction may have non-zero components in directions orthogonal to the 4d hypersurface, which implies that the system is not invariant under a shift in the 10th direction. We therefore suspect that the EK equivalence between the BFSS model and the corresponding cQEK model does not hold *as it stands.*

We may, however, add a "mass term" $\frac{1}{2}N\,m^2\int dt\,\mathrm{tr}\,X_i^2$ to the BFSS model, and consider the corresponding cQEK model, in which the mass term forces the 10th direction to be included in the 4d hypersurface. This guarantees the translational invariance in the 10th direction except at the boundary, which should be fine again due to the UV finiteness. Therefore the EK equivalence in this case is expected to hold at any finite m, and it should hold, too, even if we send m to zero eventually.

We speculate that actually quenching is not needed since the 10th direction, in which we do not have the mass term, will extend to infinity as $N\to\infty$ anyway. This implies that the IKKT model with a small mass term in 9 directions, which should be removed after taking the large N limit, is equivalent (in the sense of Eguchi-Kawai) to the BFSS model. In particular if the IKKT model has 4 extended directions, the BFSS model should have 3 out of 9 transverse directions extended, thus breaking the SO(9) symmetry down to SO(3) spontaneously. This equivalence is somewhat reminiscent of the T-duality, which relates the BFSS model on a circle to the IKKT model with Taylor's compactifying condition [23]. Note, however, that our conjecture is different in that the two models can be *both* considered in the decompactified limit.

Concluding remarks.— In this letter we have provided the first direct confirmation of the EK equivalence for the cQEK model in the case of matrix QM or the 1d gauge theory. Whether it holds also in higher dimensions is an interesting open question. This is relevant to nonperturbative studies of Matrix String Theory [24], and it is also important in view of the difficulties in formulating supersymmetric and/or chiral gauge theories on the lattice. (See Refs. [6, 25] for recent developments.) As is clear from the discussion on the SD equations, the question is highly nontrivial in the presence of UV divergences. In Ref. [12] it is speculated that the equivalence holds if the UV divergence is at most logarithmic. A natural starting point in this direction would be to study the cQEK model for the 2d pure Yang-Mills theory, and see whether the known exact results can be reproduced.

Acknowledgments.— The authors would like to thank T. Azuma, S. Iso, H. Kawai, Y. Kitazawa, K. Nagao, K. Ohta, F. Sugino, A. Tsuchiya and K. Yoshida for valuable discussions. The work of J.N. is supported in part by Grant-in-Aid for Scientific Research (No. 14740163) from the Ministry of Education, Culture, Sports, Science and Technology.

---

[*] Electronic address: kawahara@post.kek.jp
[†] Electronic address: jnishi@post.kek.jp

[1] T. Eguchi and H. Kawai, Phys. Rev. Lett. **48**, 1063 (1982).

[2] R. Dijkgraaf and C. Vafa, `hep-th/0208048`.

[3] H. Kawai, T. Kuroki and T. Morita, Nucl. Phys. B **664**, 185 (2003).

[4] T. Banks, W. Fischler, S. H. Shenker and L. Susskind, Phys. Rev. D **55**, 5112 (1997).

[5] R. A. Janik and J. Wosiek, Acta Phys. Polon. B **32**, 2143 (2001); P. Bialas and J. Wosiek, Nucl. Phys. Proc. Suppl. **106**, 968 (2002); O. Aharony, J. Marsano, S. Minwalla and T. Wiseman, Class. Quant. Grav. **21**, 5169 (2004).

[6] D. B. Kaplan and M. Unsal, `hep-lat/0503039`.

[7] G. Bhanot, U. M. Heller and H. Neuberger, Phys. Lett. B **113**, 47 (1982).

[8] M. Okawa, Phys. Rev. Lett. **49**, 353 (1982).

[9] G. Parisi, Phys. Lett. B **112**, 463 (1982).

[10] S. R. Das and S. R. Wadia, Phys. Lett. B **117**, 228 (1982).

[11] M. Okawa, Phys. Rev. Lett. **49**, 705 (1982).

[12] D. J. Gross and Y. Kitazawa, Nucl. Phys. B **206**, 440 (1982).

[13] A. Gonzalez-Arroyo and M. Okawa, Phys. Rev. D **27**, 2397 (1983).

[14] H. Aoki, N. Ishibashi, S. Iso, H. Kawai, Y. Kitazawa and T. Tada, Nucl. Phys. B **565**, 176 (2000); J. Ambjorn, Y. M. Makeenko, J. Nishimura and R. J. Szabo, JHEP **9911**, 029 (1999); Phys. Lett. B **480**, 399 (2000); JHEP **0005**, 023 (2000).

[15] A. Gonzalez-Arroyo and C. P. Korthals Altes, Phys. Lett. B **131**, 396 (1983).

[16] R. Narayanan and H. Neuberger, Phys. Rev. Lett. **91**, 081601 (2003); Nucl. Phys. B **696**, 107 (2004); J. Kiskis, R. Narayanan and H. Neuberger, Phys. Rev. D **66**, 025019 (2002); Phys. Lett. B **574**, 65 (2003); T. D. Cohen, Phys. Rev. Lett. **93**, 201601 (2004).

[17] N. Ishibashi, H. Kawai, Y. Kitazawa and A. Tsuchiya, Nucl. Phys. B **498**, 467 (1997).

[18] T. Hotta, J. Nishimura and A. Tsuchiya, Nucl. Phys. B **545**, 543 (1999).

[19] Y. M. Makeenko, *Methods of contemporary gauge theory*, (Cambridge, UK, 2002).

[20] W. Krauth and M. Staudacher, Phys. Lett. B **435**, 350 (1998).

[21] J. Ambjorn, K. N. Anagnostopoulos, W. Bietenholz, T. Hotta and J. Nishimura, JHEP **0007**, 013 (2000); J. Ambjorn, K. N. Anagnostopoulos, W. Bietenholz, F. Hofheinz and J. Nishimura, Phys. Rev. D **65**, 086001 (2002); Z. Burda, B. Petersson and M. Wattenberg, JHEP **0503**, 058 (2005).

[22] J. Nishimura and F. Sugino, JHEP **0205**, 001 (2002); H. Kawai, S. Kawamoto, T. Kuroki, T. Matsuo and S. Shinohara, Nucl. Phys. B **647**, 153 (2002); H. Kawai, S. Kawamoto, T. Kuroki and S. Shinohara, Prog. Theor. Phys. **109**, 115 (2003).

[23] W. Taylor, Phys. Lett. B **394**, 283 (1997).

[24] L. Motl, `hep-th/9701025`; R. Dijkgraaf, E. Verlinde and H. Verlinde, Nucl. Phys. B **500**, 43 (1997).

[25] D. B. Kaplan, E. Katz and M. Unsal, JHEP **0305**, 037 (2003); F. Sugino, JHEP **0501**, 016 (2005); S. Catterall, `hep-lat/0503036`.